

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 065 277 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
03.01.2001 Bulletin 2001/01

(51) Int. Cl.⁷: C12N 15/56, C12N 9/28,
C12N 1/21, C11D 3/386
// (C12N1/21, C12R1:125)

(21) Application number: 00111911.4

(22) Date of filing: 13.06.2000

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

- Hagihara, Hiroshi,
Kao Corporation
Haga-gun, Tochigi 321-3497 (JP)
- Ozaki, Katsuya,
Kao Corporation
Haga-gun, Tochigi 321-3497 (JP)

(30) Priority: 10.06.1999 JP 16356999

(71) Applicant: Kao Corporation
Tokyo 103-8210 (JP)

(74) Representative:
VOSSIUS & PARTNER
Siebertstrasse 4
81675 München (DE)

- (72) Inventors:
- Endo, Keiji,
Kao Corporation
Haga-gun, Tochigi 321-3497 (JP)
 - Igarashi, Kazuaki,
Kao Corporation
Haga-gun, Tochigi 321-3497 (JP)
 - Hayashi, Yasuhiro,
Kao Corporation
Haga-gun, Tochigi 321-3497 (JP)

Remarks:

The sequence listing, which is published as annex to the application documents, was filed after the date of filing. The applicant has declared that it does not include matter which goes beyond the content of the application as filed.

(54) **Mutant alpha-amylases**

(57) The invention relates to a mutant α -amylase obtained by making replacement or deletion of at least one of amino acid residues such as the 167th Gln, 169th Tyr and 178th Ala in the amino acid sequence set forth in SEQ ID NO:1 in an α -amylase having said amino acid sequence, or an α -amylase having a homology of at least 70% to said amino acid sequence, a gene encoding the mutant α -amylase, a vector, transformed cells, a process for producing a mutant α -amylase, comprising culturing the transformed cells, and a detergent composition comprising the mutant α -amylase.

The mutant α -amylase of the invention has excellent properties of high resistance to chelating agents, high specific activity in an alkaline region and excellent stability to heat, and is hence useful for detergents for automatic dish washer, laundry detergents and the like.

EP 1 065 277 A1

corresponds to...
your ref.: FP-KS-0555
our ref.: E1913 EP S3

Description

[0001] The present invention relates to mutant liquefying alkaline α -amylases which have excellent heat resistance, and are particularly useful as enzymes for detergents, and genes thereof.

[0002] When an α -amylase [EC.3.2.1.1] is used as an enzyme for detergents, it has heretofore been said that a liquefying alkaline α -amylase, which can decompose starch at random and is stable to alkali and also to both chelating component and oxidation bleaching component, is preferred. However, in liquefying amylases, a calcium ion is generally important for maintaining the structure of the enzymes, and the stability thereof is lowered in the presence of a chelating agent. Besides, most of such enzymes have had the optimum pH in a neutral to weakly acidic range.

[0003] Under the foregoing circumstances, the present inventors found that enzymes produced by alkaliphilic *Bacillus* sp. KSM-K38 (FERM BP-6946) and *Bacillus* sp. KSM-K36 (FERM BP-6945) strains isolated from soil do not show the lowering of activity at all in the presence of a chelating agent at a high concentration by which deactivation is recognized in the conventional liquefying α -amylases, and have resistance to surfactants and oxidizing agents and that they have higher activity on the alkaline side compared with the conventional liquefying α -amylases and are useful as enzymes for detergents (Japanese Patent Application No. 362487/1998).

[0004] However, said enzymes exhibit inactivation at a temperature of 50°C or higher, and so the heat resistance thereof has been somewhat insufficient in view of the fact that cleaning of clothing and tableware is generally conducted at about 10 to 60°C.

[0005] It is an object of the present invention to provide an α -amylase which is a liquefying alkaline α -amylase that has high activity on the alkaline side and is stable to both chelating component and oxidation bleaching component, and has excellent heat resistance.

[0006] The present inventors have acquired various mutant enzymes as to liquefying alkaline α -amylases and investigated them. As a result, it has been found that when a mutation is introduced into a specified amino acid residue in the amino acid sequence (SEQ ID NO:1) of amylase derived from KSM-K38, the heat resistance of the enzyme is improved without losing its properties such as resistance to chelating agents and resistance to oxidizing agents and high specific activity in an alkaline region, and that the heat resistance can be further improved by combining such mutations.

[0007] According to the present invention, there is thus provided a mutant α -amylase obtained by making replacement or deletion of at least one residue of amino acid residues respectively corresponding to the 11th Tyr, 16th Glu, 49th Asn, 84th Glu, 144th Ser, 167th Gln, 169th Tyr, 178th Ala, 188th Glu, 190th Asn, 205th His and 209th Gln in the amino acid sequence set forth in SEQ ID NO:1 in an α -amylase having said amino acid sequence, or an α -amylase having a homology of at least 70% to said amino acid sequence.

[0008] According to the present invention, there is also provided a mutant α -amylase obtained by making replacement of a sequence corresponding to 11 to 100 amino acid residues from the amino terminal in the amino acid sequence set forth in SEQ ID NO:1 in an α -amylase having said amino acid sequence, or an α -amylase having a homology of at least 70% to said amino acid sequence by an amino acid sequence of another liquefying α -amylase corresponding to said sequence of the amino acid residues.

[0009] According to the present invention, there are further provided genes respectively encoding these mutant α -amylases, vectors having each of the genes, cells transformed by such a vector, and a production process of these mutant α -amylases, comprising culturing the transformed cells.

[0010] According to the present invention, there is still further provided a detergent composition comprising any one of these mutant α -amylases.

[0011] The above and other objects, features and advantages of the present invention will become apparent from the following description and the appended claims, taken in conjunction with the accompanying drawings, in which:

Fig. 1 illustrates a method for preparing a recombinant plasmid for the production of α -amylases derived from KSM-K38 and KSM-AP1378 strains.

Fig. 2 illustrates a method for introducing a mutation into an α -amylase gene derived from the KSM-38 strain.

Fig. 3 illustrates a method for replacing an N-terminal sequence of the α -amylase gene derived from the KSM-38 strain by an N-terminal region of an α -amylase gene derived from the KSM-AP1378 strain.

[0012] The mutant α -amylases according to the present invention are obtained by mutating a gene encoding a liquefying alkaline α -amylase having the amino acid sequence set forth in SEQ ID NO:1 or an amino acid sequence having a homology of at least 70% to said amino acid sequence. However, an example where heat resistance is improved by deletion and/or replacement of an amino acid has also been conducted on the conventional liquefying α -amylases. For example, an enzyme obtained by deleting residues from the 177th Arg to the 178th Gly in an enzyme derived from *B. amyloliquefaciens* (J. Biol. Chem., 264, 18933, 1989) and an enzyme obtained by replacing the 133rd His in an enzyme derived from *B. licheniformis* by Tyr (J. Biol. Chem., 265, 15481, 1990) have been reported. However, the liq-

uefying alkaline α -amylases used in the present invention have a low degree of amino acid homology with the conventional liquefying alkaline α -amylases. In these α -amylases, a site corresponding to the residues from the 177th Arg to the 178th Gly has been already deleted, and the amino acid corresponding to the 133rd His has been already Tyr. Therefore, the examples of the conventional enzymes cannot be always applied. More specifically, the mutations of the amino acid sequence for improving the heat resistance in the present invention are entirely different from the examples up to the date.

[0013] Examples of the liquefying alkaline α -amylases include an enzyme (Japanese Patent Application No. 362487/1998) derived from a *Bacillus* sp. KSM-K38 (FERM BP-6946) strain separated from soil by the present inventors and having the amino acid sequence of SEQ ID NO:1 and an enzyme (SEQ ID NO:4) (Japanese Patent Application No. 362487/1998) derived from *Bacillus* sp. KSM-K36 (FERM BP-6945) and having a homology of about 95% to the amino acid sequence of SEQ ID NO:1. Incidentally, the homology of the amino acid sequence is calculated in accordance with the Lipman-Pearson method (Science, 227, 1435, 1985).

[0014] In order to obtain the mutant α -amylase according to the present invention, a gene encoding a liquefying α -amylase is first cloned from microorganisms which produce said liquefying α -amylase. As a method therefor, a general gene recombination method may be used. For example, the method described in Japanese Patent Application Laid-Open No. 336392/1996 may be used. Examples of the gene include those set forth in SEQ ID NO:3 and SEQ ID NO:5.

[0015] A mutation is then introduced into the gene thus obtained. As a method therefor, any method may be adopted so far as it is a method of site-specific mutation commonly performed. The mutation can be performed, for example, by using a Site-Directed Mutagenesis System Mutan-Super Express Km kit produced by Takara Shuzo Co., Ltd. An optional sequence of the gene may be replaced by a sequence of another gene corresponding to the optional sequence by using the recombinant PCR (polymer chain reaction) method (PCR protocols, Academic Press, New York, 1990).

[0016] The mutation for improving the heat resistance in the present invention is desirably a mutation in which an amino acid residue corresponding to the 11th Tyr in the amino acid sequence set forth in SEQ ID NO:1 is replaced by Phe, an amino acid residue corresponding to the 16th Glu by Pro, an amino acid residue corresponding to the 49th Asn by Ser, an amino acid residue corresponding to the 84th Glu by Gln, an amino acid residue corresponding to the 144th Ser by Pro, an amino acid residue corresponding to the 167th Gln by Glu, an amino acid residue corresponding to the 169th Tyr by Lys, an amino acid residue corresponding to the 178th Ala by Gln, an amino acid residue corresponding to the 188th Glu by Asp, an amino acid residue corresponding to the 190th Asn by Phe, an amino acid residue corresponding to the 205th His by Arg, or an amino acid residue corresponding to the 209th Gln by Val.

[0017] The improvement of heat resistance can also be achieved by replacing an amino acid sequence corresponding to 11 to 100 amino acid residues from the amino terminal (Asp) in the amino acid sequence of SEQ ID NO:1 according to the present invention, preferably a sequence corresponding to amino acid residues from the 1st Asp to the 19th Gly, by an amino acid sequence of another liquefying α -amylase corresponding to said sequence of the amino acid residues.

[0018] Examples of said another liquefying α -amylase used in the replacement include an enzyme having the amino acid sequence set forth in SEQ ID NO:2. A site of its amino acid sequence corresponding to said amino acid residues from the 1st Asp to the 19th Gly is from the 1st His to the 21st Gly. The enzyme is an liquefied α -amylase derived from a *Bacillus* sp. KSM-AP1378 (FERM BP-3048) strain, and the sequence of the gene is disclosed in Japanese Patent Application Laid-Open No. 336392/1996.

[0019] In the mutant α -amylases according to the present invention, a mutation with at least two kinds of replacement or deletion selected from the replacement or deletion of the above-described various kinds of amino acid residues and the replacement of the amino acid sequences combined with each other is also effective, and mutant enzymes more improved in heat resistance can be obtained by such a combination. More specifically, examples of the combination of mutations include a combination of at least two of the replacement or deletion of the various kinds of amino acid residues, a combination of at least two of the replacement of the amino acid sequence, and a combination of at least two of the replacement or deletion of the amino acid residues and the replacement of the amino acid sequence. Preferably, at least two mutations may be suitably combined from among mutations in which an amino acid residue corresponding to the 49th Asn is replaced by Ser, an amino acid residue corresponding to the 167th Gln by Glu, an amino acid residue corresponding to the 169th Tyr by Lys, an amino acid residue corresponding to the 190th Asn by Phe, an amino acid residue corresponding to the 205th His by Arg, and an amino acid residue corresponding to the 209th Gln by Val, and a mutation in which an amino acid sequence corresponding to amino acid residues from the 1st Asp to the 19th Gly is replaced by an amino acid sequence from the 1st His to the 21st Gly in the amino acid sequence set forth in SEQ ID NO:2.

[0020] Examples of the most preferred combination include a combination of mutations in which an amino acid residue corresponding to the 49th Asn is replaced by Ser, an amino acid residue corresponding to the 167th Gln by Glu, an amino acid residue corresponding to the 169th Tyr by Lys, an amino acid residue corresponding to the 190th Asn by Phe, an amino acid residue corresponding to the 205th His by Arg, and an amino acid residue corresponding to the

209th Gln by Val, and a combination of a mutation in which an amino acid sequence corresponding to amino acid residues from the 1st Asp to the 19th Gly is replaced by an amino acid sequence from the 1st His to the 21st Gly in the amino acid sequence set forth in SEQ ID NO:2 with a mutation in which an amino acid residue corresponding to the amino acid residue corresponding to the 167th Gln by Glu, an amino acid residue corresponding to the 190th Asn by Phe, or an amino acid residue corresponding to the 209th Gln by Val.

[0021] In addition, mutations for improving other properties than the heat resistance, for example, a mutation for more enhancing resistance to oxidizing agents, in which an amino acid residue corresponding to the 107th Met is replaced by Leu, a mutation for enhancing the detergency of a laundry detergent, in which an amino acid residue corresponding to the 188th Glu is replaced by Ile, and/or the like may be combined with the above-described mutations.

[0022] The thus-obtained mutant α -amylases according to the present invention are improved in stability to heat without losing excellent properties of high resistance to chelating agents, and high specific activity in an alkaline region, and are hence useful for detergents for automatic dish washer, laundry detergents and desizing agents for fibers.

[0023] Such detergents may comprise one or more enzymes selected from debranching enzymes (for example, pullulanase, isoamylase, neopullulanase, etc.), α -glycosidases, glucoamylases, proteases, cellulases, lipases, pectinases, protopectinases, pectic acid lyases, peroxidases, laccases and catalases in addition to the above-described mutant α -amylases.

[0024] Further, surfactants such as anionic surfactants, amphoteric surfactants, nonionic surfactants and cationic surfactants, chelating agents, alkalizing agents, inorganic salts, resoiling preventives, chlorine scavengers, reducing agents, bleaching agents, fluorescent dye solubilizers, perfume bases, caking preventives, enzyme activators, antioxidants, preservatives, coloring matter, bluing agents, bleaching activators, enzyme stabilizers, phase adjusters, etc., which are commonly incorporated into the classical detergents, may be incorporated.

[0025] The detergent composition according to the present invention can be produced by combining the above-described mutant α -amylases with the publicly known detergent components described above in accordance with a method known *per se* in the art. The form of the detergent composition may be suitably selected as necessary for the end application intended, and the detergent composition may be provided in the form of, for example, liquid, powder or granules. The detergent composition according to the present invention can be used as a laundry detergent, bleaching detergent, detergent for automatic dish washer, drain cleaner, artificial tooth cleaner or the like. In particular, it can preferably be used as a laundry detergent, bleaching detergent or detergent for automatic dish washer.

[0026] The mutant α -amylases according to the present invention may be used as compositions for liquefaction and saccharification of starch and be also caused to act on starch together with one or more enzymes selected from glucoamylase, maltase, pullulanase, isoamylase, neopullulanase, etc.

[0027] The mutant α -amylases according to the present invention may also be used as desizing agent compositions for fibers by incorporating an enzyme such as pullulanase, isoamylase or neopullulanase.

EXAMPLES

Determination of amylase activity and protein content:

[0028] The amylase activity and protein content of each enzyme was determined in accordance with the following respective methods.

[0029] The determination of amylase activity was conducted by the 3,5-dinitrosalicylic acid method (DNS method). After a reaction was conducted at 50°C for 15 minutes in a reaction mixture with soluble starch contained in a 50 mM glycine buffer (pH: 10), reducing sugar formed was determined by the DNS method. With respect to the enzymatic activity, the amount of the enzyme, which forms reducing sugar corresponding to 1 μ mol of glucose for 1 minute, was defined as 1 unit.

[0030] The protein content was determined by means of a Protein Assay Kit produced by Bio-Rad Laboratories making use of bovine serum albumin as a standard.

Referential Example 1:

Screening of liquefying alkaline amylase:

[0031] Soil (about 0.5 g) was suspended in sterilized water and subjected to a heat treatment at 80°C for 15 minutes. A supernatant of the heat-treated suspension was suitably diluted with sterilized water, and the resultant dilute solution was coated on an agar medium (Medium A) for separation. Culture was then conducted at 30°C for 2 days to form colonies. Those on the peripheries of which transparent halo based on amylolysis had been formed were screened, and isolated as amylase-producing bacteria. Further, the thus-isolated bacteria were inoculated on Medium B and subjected to aerobic shaking culture at 30°C for 2 days. After the culture, the resistance performance to a chelat-

ing agent (EDTA) of a supernatant centrifugally separated was determined, and its optimum pH was further measured to screen the liquefying alkaline α -amylase-producing bacteria.

[0032] *Bacillus* sp. KSM-K38 (FERM BP-6946) and *Bacillus* sp. KSM-K36 (FERM BP-6945) strains were able to be obtained by the above-described process.

Medium A:	Trypton	1.5%
	Soyton	0.5%
	Sodium chloride	0.5%
	Colored starch	0.5%
	Agar	1.5%
	Na ₂ CO ₃ (pH 10)	0.5%

Medium B:	Trypton	1.5%
	Soyton	0.5%
	Sodium chloride	0.5%
	Soluble starch	1.0%
	Na ₂ CO ₃	0.5%
	(pH 10)	

[0033] The mycological natures of the KSM-K38 and KSM-K36 strains are shown in Table 1.

Table 1

	KSM-K36 strain	KSM-K38 strain
(a) Results of microscopic observation	Bacilli having sizes of 1.0-1.2 μ m x 2.4-5.4 μ m for K36 strain and 1.0-1.2 μ m x 1.8-3.8 μ m for K38 strain. Oval endospores (1.0-1.2 μ m x 1.2-1.4 μ m) are formed at near end or the center thereof. Having periplasmic flagella and motility. Gram staining is positive. Having no acid-fast.	
(b) Growth state on various media: Incidentally, the strains are alkaliphilic and so 0.5% sodium carbonate was added to media used in the following tests.		
•Nutrient agar plate culture	Good growth state. Form of colonies is circular. Smooth surface and rough periphery. Color of colonies is pale-ocher,	Good growth state. Form of colonies is circular. Smooth surface and smooth periphery. Color of colonies is yellowish-brown.
•Nutrient agar slant culture	Grown.	Grown.
•Nutrient broth liquid culture	Grown.	Grown.

Table 1 (continued)

	KSM-K36 strain	KSM-K38 strain
5		
•Nutrient broth gelatin stab culture	Good growth state. No gelatin liquefaction is observed.	Good growth state. No gelatin liquefaction is observed.
•Litmus milk medium	Not changed.	Not changed.
(c) Physiological nature:		
10		
•Reduction of nitrate and denitrification	Reduction of nitrate is positive. Denitrification is negative.	Reduction of nitrate is positive. Denitrification is negative.
•MR test	Failed to judge because the medium is alkaline.	Failed to judge because the medium is alkaline.
•V-P test	Negative.	Negative.
15		
•Formation of indole	Negative.	Negative.
•Formation of hydrogen sulfide	Negative.	Negative.
•Hydrolysis of starch	Negative.	Negative.
20		
•Citrate utilization	Grown on Christensen medium but not grown on Coker and Simmons media.	Grown on Christensen medium but not grown on Coker and Simmons media.
•Utilization of inorganic nitrogen source	Nitrate is utilized, but ammonium salt is not utilized.	Nitrate is utilized, but ammonium salt is not utilized.
25		
•Formation of pigment	Formation of pale-yellow pigment on King B medium.	Negative.
•Urease	Negative	Negative
30		
•Oxidase	Negative	Negative
•Catalase	Negative	Negative
•Range of growth	Temperature range for growth is 15-40°C, and optimum temperature range for growth is 30-37°C. pH range for growth is pH 8.0-11.0, and optimum pH for growth is pH 10.0-11.0.	Temperature range for growth is 15-40°C, and optimum temperature for growth is 30°C. pH range for growth is pH 9.0-11.0, and optimum pH for growth is the same as described above.
35		
•Behavior against oxygen	Aerobic.	Aerobic.
40		
•O-F test	Not grown.	Not grown.
•Sugar utilization	D-galactose, D-xylose, L-arabinose, lactose, glycerol, melibiose, ribose, D-glucose, D-mannose, maltose, sucrose, trehalose, D-mannitol, starch, raffinose and D-fructose are utilized.	
45		
•Growth on salt-containing medium	Grown at a salt concentration of 12%, but not grown at a concentration of 15%.	

Reference Example 2:

Culture of KSM-K38 and KSM-K36 strains:

[0034] The KSM-K38 or KSM-K36 strain was inoculated on the liquid medium B used in Referential Example 1 to conduct shaking culture at 30°C for 2 days. The amylase activity (at pH 8.5) of a supernatant centrifugally separated was determined. As a result, these strains had activities of 557 U and 1177 U per liter of the medium, respectively.

Referential Example 3:

Purification of liquefying alkaline amylase:

- 5 **[0035]** Ammonium sulfate was added to the resultant culture supernatant of the KSM-38 strain obtained in Referential Example 2 to 80% saturation. After stirring the resultant mixture, precipitate formed was collected and dissolved in 10 mM Tris-hydrochloride buffer (pH: 7.5) containing 2 mM CaCl_2 and dialyzed overnight against the same buffer. The dialyzate thus obtained was passed through a DEAE-Toyopearl 650M column equilibrated with the same buffer and caused to be adsorbed on the column, and the intended enzyme was eluted with the same buffer by 0-1 M gradient of sodium chloride concentration. After the active fraction was dialyzed against the same buffer, an active fraction obtained by gel filtration column chromatography was dialyzed against the above-described buffer, thereby obtaining a purified enzyme which gave a single band on both polyacrylamide gel electrophoresis (gel concentration: 10%) and sodium dodecyl sulfate (SDS) electrophoresis. Incidentally, a purified enzyme was also able to be obtained from the culture supernatant of the KSM-K36 strain in accordance with the same process as described above.

Reference Example 4:

Properties of enzyme:

20 (1) Action:

[0036] Both enzymes decompose the α -1,4-glycoside bonds of starch, amylose, amylopectin and partially decomposed products thereof and produce glucose (G1), maltose (G2), maltotriose (G3), maltotetraose (G4), maltopentaose (G5), maltohexaose (G6) and maltoheptaose (G7) from amylose. However, the enzymes do not act on pullulan.

25 (2) pH stability (Britton-Robinson's buffer):

[0037] Both enzymes exhibit a residual activity of at least 70% in a pH range of 6.5 to 11 under treatment conditions of 40°C and 30 minutes.

30 (3) Action temperature range and optimum action temperature:

[0038] Both enzymes act in a wide temperature range of 20 to 80°C and have an optimum action temperature of 50 to 60°C.

35 (4) Temperature stability:

[0039] Enzyme was incubated in a 50 mM glycine-sodium hydroxide buffer (pH: 10) at various temperature for 30 minutes and then residual enzymatic activity was measured. As a result, both enzymes showed a residual activity of at least 80% at 40°C and a residual activity of about 60% even at 45°C.

(5) Molecular weight:

[0040] Both enzymes have a molecular weight of $55,000 \pm 5,000$ as measured by sodium dodecyl sulfate polyacrylamide gel electrophoresis

(6) Isoelectric point:

[0041] Both enzymes have an isoelectric point of about 4.2 as measured by isoelectric focusing.

50 (7) Influence of surfactant:

[0042] Even when both enzymes are treated at pH 10 and 30°C for 30 minutes in a 0.1% solution of each of various surfactants such as sodium linear alkylbenzenesulfonates, sodium alkylsulfates, sodium polyoxyethylene alkylsulfates, sodium α -olefinsulfonates the sodium salts of α -sulfonated fatty acid esters, sodium alkylsulfonates, SDS, soap and Softanol, they scarcely undergo inhibition of their activities (residual activity: at least 90%).

(8) Influence of metal salt:

[0043] Both enzymes were treated at pH 10 and 30°C for 30 minutes with each of various metal salts, thereby determining the influence thereof.

- 5 [0044] The K38 strain is inhibited by 1 mM Mn^{2+} (inhibitory rate: about 75%) and somewhat inhibited by both 1 mM Sr^{2+} and Cd^{2+} (inhibitory rate: 30 to 40%).

Example 1: Cloning of liquefying α -amylase gene

- 10 [0045] A chromosome DNA extracted from cells of the KSM-K38 strain in accordance with the method by Saito & Miura (Biochim. Biophys. Acta, 72, 619, 1961) was used as a template to amplify a gene fragment (about 1.5 kb) encoding a liquefying alkaline α -amylase (hereinafter referred to as "K38AMY") having an amino acid sequence set forth in SEQ ID NO:1 by PCR making use of primers K38US (SEQ ID NO: 19) and K38DH (SEQ ID NO: 20). The thus-amplified fragment was cleaved with a restriction enzyme *Sal* I, and then inserted into a *Sal* I-*Sma* I site of an expression vector
- 15 pHSP64 (Japanese Patent Application Laid-Open No. 217781/1994), thereby preparing a recombinant plasmid pHSP-K38 with a structural gene of K38AMY bonded to a trailing end of a potent promoter derived from the alkaline cellulase gene of a *Bacillus* sp. KSM-64 (FERM P-10482) strain contained in pHSP64 (Fig. 1).

- [0046] Similarly, a gene fragment (about 1.5 kb) encoding a liquefying alkaline α -amylase (hereinafter referred to as "LAMY") having an amino acid sequence set forth in SEQ ID NO:2, which had been obtained by using a chromo-
- 20 some DNA extracted from cells of a *Bacillus* sp. KSM-AP1378 (FERM BP-3048) strain (Japanese Patent Application Laid-Open No. 336392/1998) as a template, and amplified by PCR making use of primers LAUS (SEQ ID NO: 21) and LADH (SEQ ID NO: 22) was inserted into a *Sal* I-*Sma* I site of an expression vector pHSP64 in the same manner as described above, thereby preparing a recombinant plasmid pHSP-LAMY (Fig. 1).

25 Example 2: Preparation of mutant K38AMY gene-1

- [0047] A Site-Directed Mutagenesis System Mutan-Super Express Km Kit produced by Takara Shuzo Co., Ltd. was used for a site-specific mutation. The recombinant plasmid pHSP-K38 obtained in Example 1 was first used as a tem-
- 30 plate to conduct PCR making use of primers CLUBG (SEQ ID NO: 23) and K38DH (SEQ ID NO: 20), thereby amplifying a fragment of about 2.1 kb from the leading end of a potent promoter derived from the KSM-64 strain to the trailing end of the liquefying alkaline α -amylase gene. This fragment was inserted into a *Sma* I site of a plasmid vector pKF19k attached to the above kit to prepare a recombinant plasmid pKF19-K38 for introduction of mutation (Fig. 2).

- [0048] After various kinds of oligonucleotide primers for introduction of site-specific mutation respectively set forth in SEQ ID NO:6 to NO:15 were 5'-phosphorylated with a T4 DNA kinase, each of the resultant products and pKF19-
- 35 K38 were used to conduct a mutation-introducing reaction in accordance with a method described in the kit, and an *Escherichia coli* MV1184 strain (Competent Cell MV1184, product of Takara Shuzo Co., Ltd.) was transformed with the resultant reaction product. Recombinant plasmids were extracted from the resultant transformants to conduct base sequence analysis, thereby confirming the mutation.

- [0049] The mutation-introduced gene was made a template plasmid upon introduction of a different mutation by inserting an expression promoter region and a mutant K38AMY gene portion into the *Sma* I site of pKF19k again, thereby introducing another mutation in accordance with the same process as described above.

- [0050] Each of the thus-obtained mutant recombinant plasmids was used as a template to conduct PCR making use of primers CLUBG (SEQ ID NO: 23) and K38DH (SEQ ID NO: 20), thereby amplifying each of mutant K38AMY gene fragments. This fragment was cleaved with a *Sal* I and then inserted into a *Sal* I-*Sma* I site of an expression vector
- 45 pHSP64 (Japanese Patent Application Laid-Open No. 217781/1994) to prepare a plasmid for production of mutant K38AMY (Fig. 1).

Example 3: Preparation of mutant K38AMY gene-2 (chimera with LAMY gene)

- 50 [0051] Recombinant PCR was used for a mutation in which the N-terminal region of the K38AMY gene is replaced by its corresponding region of an LAMY gene (Fig. 3). The recombinant plasmid pHSP-K38 obtained in Example 1 was first used as a template to conduct PCR making use of primers K38DH (SEQ ID NO: 20) and LA-K38 (SEQ ID NO: 17), thereby amplifying a fragment encoding a sequence from the 20th Gln to C-terminal of the amino acid sequence of K38AMY set forth in SEQ ID NO: 1. On the other hand, the recombinant plasmid pHSP-LAMY was used as a template
- 55 to conduct PCR making use of primers CLUBG (SEQ ID NO: 23) and LA-K38R (SEQ ID NO: 18), thereby amplifying a gene fragment encoding a sequence from the leading end of the potent promoter to the 21st Gly of the amino acid sequence of LAMY set forth in SEQ ID NO: 2. Second PCR making use of both DNA fragments, and primers CLUBG (SEQ ID NO: 23) and K38DH (SEQ ID NO: 20) was conducted, thereby amplifying a gene fragment (about 2.1 kb)

encoding a substituted mutant enzyme (hereinafter abbreviated as "LA-K38AMY") in which both fragments having respective complementary sequences derived from the primers LA-K38 (SEQ ID NO: 17) and LA-K38R (SEQ ID NO: 18) were bonded to the terminal, and a region encoding a sequence from the 1st His to the 21st Gly of LAMY and successively a region encoding a sequence from the 20th Gln to the C-terminal of K38AMY were bonded to the trailing end of the potent promoter. This gene fragment was cleaved with *Sal* I and inserted into a *Sal* I-*Sma* I site of an expression vector pHSP64 (Japanese Patent Application Laid-Open No. 217781/1994), thereby preparing a plasmid for production of mutant K38AMY (Fig. 1).

Example 4: Production of mutant liquefying alkaline α -amylase

[0052] Each of the various plasmids for production of mutant K38AMY obtained in Examples 2 and 3 was introduced into a *Bacillus subtilis* ISW 1214 strain (*leuA metB5 hsdM1*) in accordance with the protoplast method (Mol. Gen. Genet., 168, 111, 1979) to culture the resultant recombinant *Bacillus subtilis* at 30°C for 3 days in a liquid medium (containing 8% of corn steep liquor; 1% of meat extract; 0.02% of potassium primary phosphate; 5% of maltose; 5 mM of calcium chloride; and 15 μ g/mL of tetracycline). The resultant culture supernatant was dialyzed against a Tris-HCl buffer (pH: 7.0), and the dialyzate was caused to be adsorbed on a DEAE-Toyopearl 650M column equilibrated with the same buffer, and eluted by gradient of NaCl concentration. This eluate was dialyzed against a 10 mM glycine buffer (pH: 10.0), thereby obtaining a purified enzyme of each mutant K38AMY.

Example 5: Assay of heat resistance-1

[0053] Purified preparations of an enzyme (abbreviated as "Y11F") with the 11th Tyr in SEQ ID NO:1 replaced by Phe, an enzyme (abbreviated as "N49S") with the 49th Asn replaced by Ser, an enzyme (abbreviated as "E84Q") with the 84th Glu replaced by Gln, an enzyme (abbreviated as "S144P") with the 144th Ser replaced by Pro, an enzyme (abbreviated as "Q167E") with the 167th Gln replaced by Glu, an enzyme (abbreviated as "Y169K") with the 169th Tyr replaced by Lys, an enzyme (abbreviated as "A178Q") with the 178th Ala replaced by Gln, an enzyme (abbreviated as "E188D") with the 188th Glu replaced by Asp, an enzyme (abbreviated as "N190F") with the 190th Asn replaced by Phe, and an enzyme (abbreviated as "Q209V") with the 209th Gln replaced by Val were obtained in accordance with the processes described in Examples 1, 2 and 4, and their heat resistance was assayed by the following method. As a control, wild type K38AMY was used.

[0054] Each enzyme was added to a 50 mM glycine buffer (pH: 10.0) preincubated at 50°C so as to give a concentration of about 1.2 U/mL, and after 30 minutes, the buffer was sampled to determine the residual amylase activity of the enzyme in accordance with the method described above in EXAMPLES. The activity of the enzyme at the start is regarded as 100% to determine a relative activity, thereby regarding it as the residual amylase activity. The results are shown in Table 2. In the wild type K38AMY, the residual activity was decreased to 15%, while all the mutant enzymes exhibited a high residual activity compared with the wild type.

Table 2

Enzyme	Residual activity (%) after 30 minutes
Wild type	15
Y11F	40
N49S	30
E84Q	25
S144P	30
Q167E	46
Y169K	63
A178Q	20
E188D	30
N190F	70
Q209V	40

Example 6: Assay of heat resistance-2

[0055] Mutant enzymes with Q167E, Y169K, N190F and Q209V among the mutations described in Example 5 combined in the following manner were prepared in accordance with the processes described in Examples 1, 2 and 4.

Q167E/Y169K (abbreviated as "QEYK", prepared by using primer of SEQ ID NO: 16)

N190F/Q209V (abbreviated as "NFQV")

Q167E/Y169K/N190F/Q209V (abbreviated as "QEYK/NFQV")

[0056] With respect to these enzymes, the heat resistance was assayed by a method similar to Example 5. However, the temperature in the heat treatment was changed to 55°C, and Q167E, Y169K, N190F and Q209V were used as controls. As a result, as shown in Table 3, all the mutants were observed being improved in heat resistance by the combination, and QEYK/NFQV obtained by combining 4 mutations exhibited a residual activity of 85% after 30 minutes even at 55°C.

Table 3

Enzyme	Residual activity (%) after 30 minutes
Q167E	7
Y169K	14
QEYK	45
N190F	20
Q209V	1
NFQV	40
QEYK/NFQV	85

Example 7: Assay of heat resistance-3

[0057] The following mutant enzymes with the mutation NFQV described in Example 6 combined with S144P described in Example 5, and further combined with a mutation of replacement of 16th Gln by Pro (abbreviated as "E16P") were prepared in accordance with the processes described in Examples 1, 2 and 4.

S144P/NFQV (abbreviated as "SP/NFQV")

E16P/S144P/NFQV (abbreviated as "EPSP/NFQV")

[0058] With respect to these enzymes, the heat resistance was assayed by a method (50°C) similar to Example 5. As a result, as shown in Table 4, improvement in heat resistance was observed by combining E16P with SP/NFQV.

Table 4

Enzyme	Residual activity (%) after 30 minutes
SP/NFQV	40
EPSP/NFQV	50

Example 8: Assay of heat resistance-4

[0059] The following mutant enzymes with QEYK/NFQV among the mutations described in Example 6 suitably combined with a mutation (abbreviated as "M107L") with the 107th Met in SEQ ID NO:1 replaced by Leu, a mutation (abbreviated as "H205R") with the 205th His replaced by Arg, and N49S among the mutations described in Example 5 were prepared in accordance with the processes described in Examples 1, 2 and 4.

M107L/QEYK/NFQV (abbreviated as "ML/QEYK/NFQV")

N49S/M107L/QEYK/NFQV (abbreviated as "NSML/QEYK/NFQV")

N49S/M107L/H205R/QEYK/NFQV (abbreviated as "NSMLHR/QEYK/NFQV")

5 [0060] With respect to these enzymes, the heat resistance was assayed by a method similar to Example 5. However, the temperature in the heat treatment was changed to 60°C.

[0061] As a result, heat resistance was additionally improved by combining ML/QEYK/NFQV with N49S, further H205R, and NSMLHR/QEYK/NFQV exhibited a residual activity of 75% after 30 minutes even at 60°C (Table 5)

Table 5

Enzyme	Residual activity (%) after 30 minutes
ML/QEYK/NFQV	30
NSML/QEYK/NFQV	50
NSMLHR/QEYK/NFQV	75

20 Example 9: Assay of heat resistance-5

[0062] A mutant enzyme LA-K38AMY with a sequence from the 1st Asp to the 19th Gly of K38AMY replaced by a sequence from the 1st His to the 21st Gly of LAMY was obtained in accordance with the processes described in Examples 1, 3 and 4. The heat resistance of this enzyme was assayed by the method described in Example 5. As a result, as shown in Table 6, improvement in heat resistance by the replacement was observed.

Table 6

Enzyme	Residual activity (%) after 30 minutes
Wild type	15
LA-K38AMY	33

35 Example 10: Assay of heat resistance-6

[0063] Into the gene of the mutant enzyme QEYK/NFQV described in Example 6, was introduced a mutation with a sequence from the 1st Asp to the 19th Gly replaced by a sequence from the 1st His to the 21st Gly of LAMY in accordance with the same processes as in Examples 1 and 3. With respect to a mutant enzyme LA-K38AMY/QEYK/NFQV obtained by using this enzyme in accordance with the process described in Example 4, the heat resistance was assayed by the same method (heat treatment temperature: 60°C) as in Example 8.

[0064] As a result, heat resistance was additionally improved by the combination, and LA-K38AMY/QEYK/NFQV exhibited a residual activity of 63% after 30 minutes even at 60°C (Table 7)

Table 7

Enzyme	Residual activity (%) after 30 minutes
LA-K38AMY	1
QEYK/NFQV	40
LA-K38AMY/QEYK/NFQV	63

55 Example 11: Detergent composition for automatic dish washer

[0065] A detergent composition for automatic dish washer was produced in accordance with a formulation shown

in Table 8, and various mutant enzymes were separately incorporated into this detergent composition to conduct a washing test. As a result, the mutant enzymes exhibited an excellent detergent effect compared with the wild type enzyme when the enzymes having the same activity value as each other are added.

Table 8

Composition of detergent	(%)
Pluronic L-61	2.2
Sodium carbonate	24.7
Sodium hydrogencarbonate	24.7
Sodium percarbonate	10.0
Sodium silicate No. 1	12.0
Trisodium citrate	20.0
Polypropylene glycol	2.2
Silicone KST-04 (product of Toshiba silicone Co., Ltd.)	0.2
Socarane CP-A45 (product of BASF AG)	4.0

[0066] The mutant α -amylases according to the present invention have excellent properties of high resistance to chelating agents, high specific activity in an alkaline region, excellent stability to heat, and are hence useful for detergents for automatic dish washer, laundry detergents, compositions for liquefaction and saccharification of starch, and desizing agents for fibers.

<110> KAO CORPORATION

<130> E 1913 EP

<140> 00 11 1911.4

<141> 2000-06-13

<150>JP P1999-163569

<151>1999-06-10

<160>23

<210>1

<211>480

<212>PRT

<213>Bacillus sp. KSM-K38

<400>1

Asp Gly Leu Asn Gly Thr Met Met Gln Tyr Tyr Glu Trp His Leu Glu
5 10 15

Asn Asp Gly Gln His Trp Asn Arg Leu His Asp Asp Ala Ala Ala Leu

20 25 30

Ser Asp Ala Gly Ile Thr Ala Ile Trp Ile Pro Pro Ala Tyr Lys Gly

35	40	45
----	----	----

Asn Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu

50 55 60

Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
65 70 75 80

65 70 75 80
Ala Glu Leu Gly Arg Ala Ile Gly Ser Leu Lys Ser Asp Asp Ile Asp

Ala Gln Leu Glu Arg Ala Ile Gly Ser Leu Lys Ser Asn Asp Ile Asn
85 90 95

Val Tyr Gly Asn Val Val Met Asn His Lys Met Gly Ala Asp Phe Thr

Val	Ile	Gly	Asp	Val	Val	Met	Asn	His	Gly	Met	Gly	Leu	Asp	Leu	Leu
			100					105					110		

Glu Ala Val Gln Ala Val Gln Val Asn Pro Thr Asn Arg Trp Gln Asp

115	120	125
-----	-----	-----

Ile Ser Gly Ala Tyr Thr Ile Asp Ala Trp Thr Gly Phe Asp Phe Ser

130 135 140

Gly Arg Asn Asn Ala Tyr Ser Asp Phe Lys Trp Arg Trp Phe His Phe

145	150	155	160
Jan	Feb	Mar	Apr
May	Jun	Jul	Aug
Sep	Oct	Nov	Dec

Asn Gly Val Asp Trp Asp Gln Arg Tyr Gln Glu Asn His Ile Phe Arg
165 170 175

165 170 175
Phe Ala Asp Thr Asp Trp Asp Trp Arg Val Asp Glu Glu Asp Gly Asn

Phe Ala Asn Thr Asn Trp Asn Trp Arg Val Asp Glu Glu Asn Gly Asn
180 185 190

180 185 190
Tyr Asp Tyr Leu Leu Gly Ser Asn Ile Asp Phe Ser His Pro Glu Val

[illegible]

Gln Asp Glu Leu Lys Asp Trp Gly Ser Trp Phe Thr Asp Glu Leu Asp

210 215 220

Leu Asp Gly Tyr Arg Leu Asp Ala Ile Lys His Ile Pro Phe Trp Tyr

225 230 235 240

Thr Ser Asp Trp Val Arg His Gln Arg Asn Glu Ala Asp Gln Asp Leu

245 250 255
 Gls Hal Hal Gls Gls Ten Ten Ten Ten Hal Gls Gls Ten Gls Ph

Phe Val Val Gly Glu Tyr Trp Lys Asp Asp Val Gly Ala Leu Glu Phe
360 365 370

260			265			270									
Trp	Leu	Asp	Gly	Met	Asp	Trp	Gly	Met	Ser	Leu	Phe	Asp	Val	Pro	Leu
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96
97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112
113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128
129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144
145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176
177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192
193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208
209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224
225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240
241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256
257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272
273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288
289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304
305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320
321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336
337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352
353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368
369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384
385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400
401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416
417	418	419	420	421	422	423	424	425	426						

Tyr Leu Asp Glu Met Asn Trp Glu Met Ser Leu Phe Asp Val Pro Leu
275 280 285

Asn Tyr Asn Phe Tyr Arg Ala Ser Gln Gln Gly Gly Ser Tyr Asp Met

ASN	TY1	ASN	PHE	TY1	ARG	ALA	SER	GLN	GLN	GLY	GLY	SER	TY1	ASP	ASP
290						295					300				

Arg Asn Ile Leu Arg Gly Ser Leu Val Glu Ala His Pro Met His Ala
 305 310 315 320
 Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Glu Ser Leu Glu
 325 330 335
 Ser Trp Val Ala Asp Trp Phe Lys Pro Leu Ala Tyr Ala Thr Ile Leu
 340 345 350
 Thr Arg Glu Gly Gly Tyr Pro Asn Val Phe Tyr Gly Asp Tyr Tyr Gly
 355 360 365
 Ile Pro Asn Asp Asn Ile Ser Ala Lys Lys Asp Met Ile Asp Glu Leu
 370 375 380
 Leu Asp Ala Arg Gln Asn Tyr Ala Tyr Gly Thr Gln His Asp Tyr Phe
 385 390 395 400
 Asp His Trp Asp Val Gly Trp Thr Arg Glu Gly Ser Ser Ser Arg
 405 410 415
 Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asn Gly Pro Gly Gly Ser
 420 425 430
 Lys Trp Met Tyr Val Gly Arg Gln Asn Ala Gly Gln Thr Trp Thr Asp
 435 440 445
 Leu Thr Gly Asn Asn Gly Ala Ser Val Thr Ile Asn Gly Asp Gly Trp
 450 455 460
 Gly Glu Phe Phe Thr Asn Gly Gly Ser Val Ser Val Tyr Val Asn Gln
 465 470 475 480

<210>2

<211>485

<212>PRT

<213>Bacillus sp. KSM-AP1378

<400>2

His His Asn Gly Thr Asn Gly Thr Met Met Gln Tyr Phe Glu Trp His
 5 10 15
 Leu Pro Asn Asp Gly Asn His Trp Asn Arg Leu Arg Asp Asp Ala Ala
 20 25 30
 Asn Leu Lys Ser Lys Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Trp
 35 40 45
 Lys Gly Thr Ser Gln Asn Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr
 50 55 60
 Asp Leu Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly
 65 70 75 80
 Thr Arg Ser Gln Leu Gln Gly Ala Val Thr Ser Leu Lys Asn Asn Gly
 85 90 95
 Ile Gln Val Tyr Gly Asp Val Val Met Asn His Lys Gly Gly Ala Asp
 100 105 110
 Gly Thr Glu Met Val Asn Ala Val Glu Val Asn Arg Ser Asn Arg Asn
 115 120 125
 Gln Glu Ile Ser Gly Glu Tyr Thr Ile Glu Ala Trp Thr Lys Phe Asp
 130 135 140
 Phe Pro Gly Arg Gly Asn Thr His Ser Asn Phe Lys Trp Arg Trp Tyr
 145 150 155 160
 His Phe Asp Gly Thr Asp Trp Asp Gln Ser Arg Gln Leu Gln Asn Lys
 165 170 175
 Ile Tyr Lys Phe Arg Gly Thr Gly Lys Ala Trp Asp Trp Glu Val Asp
 180 185 190
 Ile Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Ile Asp Met
 195 200 205
 Asp His Pro Glu Val Ile Asn Glu Leu Arg Asn Trp Gly Val Trp Tyr
 210 215 220
 Thr Asn Thr Leu Asn Leu Asp Gly Phe Arg Ile Asp Ala Val Lys His
 225 230 235 240
 Ile Lys Tyr Ser Tyr Thr Arg Asp Trp Leu Thr His Val Arg Asn Thr

5 Thr Gly Lys Pro Met Phe Ala Val Ala Glu Phe Trp Lys Asn Asp Leu
 245 250 255
 260 265 270
 Ala Ala Ile Glu Asn Tyr Leu Asn Lys Thr Ser Trp Asn His Ser Val
 275 280 285
 Phe Asp Val Pro Leu His Tyr Asn Leu Tyr Asn Ala Ser Asn Ser Gly
 290 295 300
 Gly Tyr Phe Asp Met Arg Asn Ile Leu Asn Gly Ser Val Val Gln Lys
 305 310 315 320
 10 His Pro Ile His Ala Val Thr Phe Val Asp Asn His Asp Ser Gln Pro
 325 330 335
 Gly Glu Ala Leu Glu Ser Phe Val Gln Ser Trp Phe Lys Pro Leu Ala
 340 345 350
 Tyr Ala Leu Ile Leu Thr Arg Glu Gln Gly Tyr Pro Ser Val Phe Tyr
 355 360 365
 15 Gly Asp Tyr Tyr Gly Ile Pro Thr His Gly Val Pro Ser Met Lys Ser
 370 375 380
 Lys Ile Asp Pro Leu Leu Gln Ala Arg Gln Thr Tyr Ala Tyr Gly Thr
 385 390 395 400
 Gln His Asp Tyr Phe Asp His His Asp Ile Ile Gly Trp Thr Arg Glu
 405 410 415
 20 Gly Asp Ser Ser His Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asp
 420 425 430
 Gly Pro Gly Gly Asn Lys Trp Met Tyr Val Gly Lys His Lys Ala Gly
 435 440 445
 Gln Val Trp Arg Asp Ile Thr Gly Asn Arg Ser Gly Thr Val Thr Ile
 450 455 460
 25 Asn Ala Asp Gly Trp Gly Asn Phe Thr Val Asn Gly Gly Ala Val Ser
 465 470 475 480
 Val Trp Val Lys Gln
 485

 <210>3
 30 <211>1753
 <212>DNA
 <213>Bacillus sp. KSM-K38

 <220>
 <221>sig_peptide
 35 <222>(162)..(224)

 <220>
 <221>mat_peptide
 <222>(225)..(1664)

 40 <220>
 <221>CDS
 <222>(162)..(1664)

 <400>3
 45 gtagcgaaa cgatgcgcaa aactgcgcaa ctactagcac tcttcagggga ctaaaccacc 60
 ttttttccaa aaatgacatc atataaaca atttgtctac caatcactat ttaaagctgt 120
 ttatgatata tgtaagcgtt atcattaataa ggaggtattt g atg aga aga tgg gta 176
 gta gca atg ttg gca gtg tta ttt tta ttt cct tcg gta gta gtt gca 224
 gat gga ttg aac ggt acg atg atg cag tat tat gag tgg cat ttg gaa 272
 Asp Gly Leu Asn Gly Thr Met Met Gln Tyr Tyr Glu Trp His Leu Glu
 1 5 10 15
 50 aac gac ggg cag cat tgg aat cgg ttg cac gat gat gcc gca gct ttg 320
 Asn Asp Gly Gln His Trp Asn Arg Leu His Asp Asp Ala Ala Leu
 20 25 30

agt gat gct ggt att aca gct att tgg att ccg cca gcc tac aaa ggt 368
 Ser Asp Ala Gly Ile Thr Ala Ile Trp Ile Pro Pro Ala Tyr Lys Gly
 35 40 45
 5 aat agt cag gcg gat gtt ggg tac ggt gca tac gat ctt tat gat tta 416
 Asn Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu
 50 55 60
 gga gag ttc aat caa aag ggt act gtt cga acg aaa tac gga act aag 464
 Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
 65 70 75 80
 10 gca cag ctt gaa cga gct att ggg tcc ctt aaa tct aat gat atc aat 512
 Ala Gln Leu Glu Arg Ala Ile Gly Ser Leu Lys Ser Asn Asp Ile Asn
 85 90 95
 gta tac gga gat gtc gtg atg aat cat aaa atg gga gct gat ttt acg 560
 Val Tyr Gly Asp Val Val Met Asn His Lys Met Gly Ala Asp Phe Thr
 100 105 110
 15 gag gca gtg caa gct gtt caa gta aat cca acg aat cgt tgg cag gat 608
 Glu Ala Val Gln Ala Val Gln Val Asn Pro Thr Asn Arg Trp Gln Asp
 115 120 125
 att tca ggt gcc tac acg att gat gcg tgg acg ggt ttc gac ttt tca 656
 Ile Ser Gly Ala Tyr Thr Ile Asp Ala Trp Thr Gly Phe Asp Phe Ser
 130 135 140
 20 ggg cgt aac aac gcc tat tca gat ttt aag tgg aga tgg ttc cat ttt 704
 Gly Arg Asn Asn Ala Tyr Ser Asp Phe Lys Trp Arg Trp Phe His Phe
 145 150 155 160
 aat ggt gtt gac tgg gat cag cgc tat caa gaa aat cat att ttc cgc 752
 Asn Gly Val Asp Trp Asp Gln Arg Tyr Gln Glu Asn His Ile Phe Arg
 165 170 175
 25 ttt gca aat acg aac tgg aac tgg cga gtg gat gaa gag aac ggt aat 800
 Phe Ala Asn Thr Asn Trp Asn Trp Arg Val Asp Glu Glu Asn Gly Asn
 180 185 190
 tat gat tac ctg tta gga tcg aat atc gac ttt agt cat cca gaa gta 848
 Tyr Asp Tyr Leu Leu Gly Ser Asn Ile Asp Phe Ser His Pro Glu Val
 195 200 205
 30 caa gat gag ttg aag gat tgg ggt agc tgg ttt acc gat gag tta gat 896
 Gln Asp Glu Leu Lys Asp Trp Gly Ser Trp Phe Thr Asp Glu Leu Asp
 210 215 220
 ttg gat ggt tat cgt tta gat gct att aaa cat att cca ttc tgg tat 944
 Leu Asp Gly Tyr Arg Leu Asp Ala Ile Lys His Ile Pro Phe Trp Tyr
 225 230 235 240
 35 aca tct gat tgg gtt cgg cat cag cgc aac gaa gca gat caa gat tta 992
 Thr Ser Asp Trp Val Arg His Gln Arg Asn Glu Ala Asp Gln Asp Leu
 245 250 255
 ttt gtc gta ggg gaa tat tgg aag gat gac gta ggt gct ctc gaa ttt 1040
 Phe Val Val Gly Glu Tyr Trp Lys Asp Asp Val Gly Ala Leu Glu Phe
 260 265 270
 40 tat tta gat gaa atg aat tgg gag atg tct cta ttc gat gtt cca ctt 1088
 Tyr Leu Asp Glu Met Asn Trp Glu Met Ser Leu Phe Asp Val Pro Leu
 275 280 285
 aat tat aat ttt tac cgg gct tca caa caa ggt gga agc tat gat atg 1136
 Asn Tyr Asn Phe Tyr Arg Ala Ser Gln Gln Gly Gly Ser Tyr Asp Met
 290 295 300
 45 cgt aat att tta cga gga tct tta gta gaa gcg cat ccg atg cat gca 1184
 Arg Asn Ile Leu Arg Gly Ser Leu Val Glu Ala His Pro Met His Ala
 305 310 315 320
 gtt acg ttt gtt gat aat cat gat act cag cca ggg gag tca tta gag 1232
 Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Glu Ser Leu Glu
 325 330 335
 50 tca tgg gtt gct gat tgg ttt aag cca ctt gct tat gcg aca att ttg 1280
 Ser Trp Val Ala Asp Trp Phe Lys Pro Leu Ala Tyr Ala Thr Ile Leu
 340 345 350

55

acg cgt gaa ggt ggt tat cca aat gta ttt tac ggt gat tac tat ggg 1328
 Thr Arg Glu Gly Gly Tyr Pro Asn Val Phe Tyr Gly Asp Tyr Tyr Gly
 355 360 365
 5 att cct aac gat aac att tca gct aaa aaa gat atg att gat gag ctg 1376
 Ile Pro Asn Asp Asn Ile Ser Ala Lys Lys Asp Met Ile Asp Glu Leu
 370 375 380
 ctt gat gca cgt caa aat tac gca tat ggc acg cag cat gac tat ttt 1424
 Leu Asp Ala Arg Gln Asn Tyr Ala Tyr Gly Thr Gln His Asp Tyr Phe
 385 390 395 400
 10 gat cat tgg gat gtt gta gga tgg act agg gaa gga tct tcc tcc aga 1472
 Asp His Trp Asp Val Val Gly Trp Thr Arg Glu Gly Ser Ser Ser Arg
 405 410 415
 cct aat tca ggc ctt gcg act att atg tgc aat gga cct ggt ggt tcc 1520
 Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asn Gly Pro Gly Gly Ser
 420 425 430
 15 aag tgg atg tat gta gga cgt cag aat gca gga caa aca tgg aca gat 1568
 Lys Trp Met Tyr Val Gly Arg Gln Asn Ala Gly Gln Thr Trp Thr Asp
 435 440 445
 tta act ggt aat aac gga gcg tcc gtt aca att aat ggc gat gga tgg 1616
 Leu Thr Gly Asn Asn Gly Ala Ser Val Thr Ile Asn Gly Asp Gly Trp
 450 455 460
 20 ggc gaa ttc ttt acg aat gga gga tct gta tcc gtg tac gtg aac caa 1664
 Gly Glu Phe Phe Thr Asn Gly Gly Ser Val Ser Val Tyr Val Asn Gln
 465 470 475 480
 taacaaaaag ccttgagaag ggattcctcc ctaactcaag gctttcttta tgcgcttag 1724
 cttaacgctt ctacgacttt gaagcttta 1753
 25
 <210>4
 <211>480
 <212>PRT
 <213>Bacillus sp. KSM-K36
 30
 <400>4
 Asp Gly Leu Asn Gly Thr Met Met Gln Tyr Tyr Glu Trp His Leu Glu
 5 10 15
 Asn Asp Gly Gln His Trp Asn Arg Leu His Asp Asp Ala Glu Ala Leu
 20 25 30
 Ser Asn Ala Gly Ile Thr Ala Ile Trp Ile Pro Pro Ala Tyr Lys Gly
 35 40 45
 Asn Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu
 50 55 60
 Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
 65 70 75 80
 Ala Gln Leu Glu Arg Ala Ile Gly Ser Leu Lys Ser Asn Asp Ile Asn
 85 90 95
 40 Val Tyr Gly Asp Val Val Met Asn His Lys Leu Gly Ala Asp Phe Thr
 100 105 110
 Glu Ala Val Gln Ala Val Gln Val Asn Pro Ser Asn Arg Trp Gln Asp
 115 120 125
 Ile Ser Gly Val Tyr Thr Ile Asp Ala Trp Thr Gly Phe Asp Phe Pro
 130 135 140
 45 Gly Arg Asn Asn Ala Tyr Ser Asp Phe Lys Trp Arg Trp Phe His Phe
 145 150 155 160
 Asn Gly Val Asp Trp Asp Gln Arg Tyr Gln Glu Asn His Leu Phe Arg
 165 170 175
 Phe Ala Asn Thr Asn Trp Asn Trp Arg Val Asp Glu Glu Asn Gly Asn
 180 185 190
 50 Tyr Asp Tyr Leu Leu Gly Ser Asn Ile Asp Phe Ser His Pro Glu Val
 195 200 205

55

5 Gln Glu Glu Leu Lys Asp Trp Gly Ser Trp Phe Thr Asp Glu Leu Asp
 210 215 220
 Leu Asp Gly Tyr Arg Leu Asp Ala Ile Lys His Ile Pro Phe Trp Tyr
 225 230 235 240
 Thr Ser Asp Trp Val Arg His Gln Arg Ser Glu Ala Asp Gln Asp Leu
 245 250 255
 Phe Val Val Gly Glu Tyr Trp Lys Asp Asp Val Gly Ala Leu Glu Phe
 260 265 270
 Tyr Leu Asp Glu Met Asn Trp Glu Met Ser Leu Phe Asp Val Pro Leu
 275 280 285
 10 Asn Tyr Asn Phe Tyr Arg Ala Ser Lys Gln Gly Gly Ser Tyr Asp Met
 290 295 300
 Arg Asn Ile Leu Arg Gly Ser Leu Val Glu Ala His Pro Ile His Ala
 305 310 315 320
 Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Glu Ser Leu Glu
 325 330 335
 15 Ser Trp Val Ala Asp Trp Phe Lys Pro Leu Ala Tyr Ala Thr Ile Leu
 340 345 350
 Thr Arg Glu Gly Gly Tyr Pro Asn Val Phe Tyr Gly Asp Tyr Tyr Gly
 355 360 365
 Ile Pro Asn Asp Asn Ile Ser Ala Lys Lys Asp Met Ile Asp Glu Leu
 370 375 380
 20 Leu Asp Ala Arg Gln Asn Tyr Ala Tyr Gly Thr Gln His Asp Tyr Phe
 385 390 395 400
 Asp His Trp Asp Ile Val Gly Trp Thr Arg Glu Gly Thr Ser Ser Arg
 405 410 415
 Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asn Gly Pro Gly Gly Ser
 420 425 430
 25 Lys Trp Met Tyr Val Gly Gln Gln His Ala Gly Gln Thr Trp Thr Asp
 435 440 445
 Leu Thr Gly Asn His Ala Ala Ser Val Thr Ile Asn Gly Asp Gly Trp
 450 455 460
 Gly Glu Phe Phe Thr Asn Gly Gly Ser Val Ser Val Tyr Val Asn Gln
 465 470 475 480

 <210>5
 <211>1625
 <212>DNA
 <213>Bacillus sp.KSM-K36

 35 <220>
 <221>sig_peptide
 <222>(40)..(102)

 <220>
 <221>mat_peptide
 40 <222>(103)..(1542)

 <220>
 <221>CDS
 <222>(40)..(1542)

 45 <400>5
 atgatatatg taagcggttat cattaaaagg aggtatttg atg aaa aga tgg gta 54
 gta gca atg ctg gca gtg tta ttt tta ttt cct tcg gta gta gtt gca 102
 gat ggc ttg aat gga acg atg atg cag tat tat gag tgg cat cta gag 150
 Asp Gly Leu Asn Gly Thr Met Met Gln Tyr Tyr Glu Trp His Leu Glu
 1 5 10 15
 50 aat gat ggg caa cac tgg aat cgg ttg cat gat gat gcc gaa gct tta 198
 Asn Asp Gly Gln His Trp Asn Arg Leu His Asp Asp Ala Glu Ala Leu

				20				25					30				
	agt	aat	gcg	ggg	att	aca	gct	att	tgg	ata	ccc	cca	gcc	tac	aaa	gga	246
5	Ser	Asn	Ala	Gly	Ile	Thr	Ala	Ile	Trp	Ile	Pro	Pro	Ala	Tyr	Lys	Gly	
		35						40					45				
	aat	agt	cag	gct	gat	gtt	ggg	tat	ggg	gca	tac	gac	ctt	tat	gat	tta	294
	Asn	Ser	Gln	Ala	Asp	Val	Gly	Tyr	Gly	Ala	Tyr	Asp	Leu	Tyr	Asp	Leu	
		50					55					60					
	ggg	gag	ttt	aat	caa	aaa	ggg	acc	gtt	cga	acg	aaa	tac	ggg	aca	aag	342
10	Gly	Glu	Phe	Asn	Gln	Lys	Gly	Thr	Val	Arg	Thr	Lys	Tyr	Gly	Thr	Lys	
		65				70					75					80	
	gct	cag	ctt	gag	cga	gct	ata	ggg	tcc	cta	aag	tcg	aat	gat	atc	aat	390
	Ala	Gln	Leu	Glu	Arg	Ala	Ile	Gly	Ser	Leu	Lys	Ser	Asn	Asp	Ile	Asn	
					85					90					95		
	gtt	tat	ggg	gat	gtc	gta	atg	aat	cat	aaa	tta	gga	gct	gat	ttc	acg	438
15	Val	Tyr	Gly	Asp	Val	Val	Met	Asn	His	Lys	Leu	Gly	Ala	Asp	Phe	Thr	
				100				105						110			
	gag	gca	gtg	caa	gct	gtt	caa	gta	aat	cct	tcg	aac	cgt	tgg	cag	gat	486
	Glu	Ala	Val	Gln	Ala	Val	Gln	Val	Asn	Pro	Ser	Asn	Arg	Trp	Gln	Asp	
				115				120					125				
	att	tca	ggg	gtc	tac	acg	att	gat	gca	tgg	acg	gga	ttt	gac	ttt	cca	534
20	Ile	Ser	Gly	Val	Tyr	Thr	Ile	Asp	Ala	Trp	Thr	Gly	Phe	Asp	Phe	Pro	
				130				135					140				
	ggg	cgc	aac	aat	gcc	tat	tcc	gat	ttt	aaa	tgg	aga	tgg	ttc	cat	ttt	582
	Gly	Arg	Asn	Asn	Ala	Tyr	Ser	Asp	Phe	Lys	Trp	Arg	Trp	Phe	His	Phe	
						150					155					160	
	aat	ggc	gtt	gac	tgg	gat	caa	cgc	tat	caa	gaa	aac	cat	ctt	ttt	cgc	630
25	Asn	Gly	Val	Asp	Trp	Asp	Gln	Arg	Tyr	Gln	Glu	Asn	His	Leu	Phe	Arg	
					165					170						175	
	ttt	gca	aat	acg	aac	tgg	aac	tgg	cga	gtg	gat	gaa	gag	aat	ggg	aat	678
	Phe	Ala	Asn	Thr	Asn	Trp	Asn	Trp	Arg	Val	Asp	Glu	Glu	Asn	Gly	Asn	
					180				185					190			
	tat	gac	tat	tta	tta	gga	tcg	aac	att	gac	ttt	agc	cac	cca	gag	gtt	726
30	Tyr	Asp	Tyr	Leu	Leu	Gly	Ser	Asn	Ile	Asp	Phe	Ser	His	Pro	Glu	Val	
						195		200					205				
	caa	gag	gaa	tta	aag	gat	tgg	ggg	agc	tgg	ttt	acg	gat	gag	cta	gat	774
	Gln	Glu	Glu	Leu	Lys	Asp	Trp	Gly	Ser	Trp	Phe	Thr	Asp	Glu	Leu	Asp	
						210		215					220				
	tta	gat	ggg	tat	cga	ttg	gat	gct	att	aag	cat	att	cca	ttc	tgg	tat	822
35	Leu	Asp	Gly	Tyr	Arg	Leu	Asp	Ala	Ile	Lys	His	Ile	Pro	Phe	Trp	Tyr	
						225		230				235				240	
	acg	tca	gat	tgg	gtt	agg	cat	cag	cga	agt	gaa	gca	gac	caa	gat	tta	870
	Thr	Ser	Asp	Trp	Val	Arg	His	Gln	Arg	Ser	Glu	Ala	Asp	Gln	Asp	Leu	
						245				250					255		
	ttt	gtc	gta	ggg	gag	tat	tgg	aag	gat	gac	gta	ggg	gct	ctc	gaa	ttt	918
40	Phe	Val	Val	Gly	Glu	Tyr	Trp	Lys	Asp	Asp	Val	Gly	Ala	Leu	Glu	Phe	
				260					265					270			
	tat	tta	gat	gaa	atg	aat	tgg	gag	atg	tct	cta	ttc	gat	gtt	ccg	ctc	966
	Tyr	Leu	Asp	Glu	Met	Asn	Trp	Glu	Met	Ser	Leu	Phe	Asp	Val	Pro	Leu	
						275			280					285			
	aat	tat	aat	ttt	tac	cgg	gct	tca	aag	caa	ggc	gga	agc	tat	gat	atg	1014
45	Asn	Tyr	Asn	Phe	Tyr	Arg	Ala	Ser	Lys	Gln	Gly	Gly	Ser	Tyr	Asp	Met	
						295						300					
	cgt	aat	att	tta	cga	gga	tct	tta	gta	gaa	gca	cat	ccg	att	cat	gca	1062
	Arg	Asn	Ile	Leu	Arg	Gly	Ser	Leu	Val	Glu	Ala	His	Pro	Ile	His	Ala	
						310						315				320	
	gtt	acg	ttt	gtt	gat	aat	cat	gat	act	cag	cca	gga	gag	tca	tta	gaa	1110
50	Val	Thr	Phe	Val	Asp	Asn	His	Asp	Thr	Gln	Pro	Gly	Glu	Ser	Leu	Glu	
						325					330					335	
	tca	tgg	gtc	gct	gat	tgg	ttt	aag	cca	ctt	gct	tat	gcg	aca	atc	ttg	1158
	Ser	Trp	Val	Ala	Asp	Trp	Phe	Lys	Pro	Leu	Ala	Tyr	Ala	Thr	Ile	Leu	

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

340
 acg cgt gaa ggt ggt tat cca aat gta ttt tac ggt gac tac tat ggg 1206
 Thr Arg Glu Gly Gly Tyr Pro Asn Val Phe Tyr Gly Asp Tyr Tyr Gly
 355 360 365
 att cct aac gat aac att tca gct aag aag gat atg att gat gag ttg 1254
 Ile Pro Asn Asp Asn Ile Ser Ala Lys Lys Asp Met Ile Asp Glu Leu
 370 375 380
 ctt gat gca cgt caa aat tac gca tat ggc aca caa cat gac tat ttt 1302
 Leu Asp Ala Arg Gln Asn Tyr Ala Tyr Gly Thr Gln His Asp Tyr Phe
 385 390 395 400
 gat cat tgg gat atc gtt gga tgg aca aga gaa ggt aca tcc tca cgt 1350
 Asp His Trp Asp Ile Val Gly Trp Thr Arg Glu Gly Thr Ser Ser Arg
 405 410 415
 cct aat tcg ggt ctt gct act att atg tcc aat ggt cct gga gga tca 1398
 Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asn Gly Pro Gly Gly Ser
 420 425 430
 aaa tgg atg tac gta gga cag caa cat gca gga caa acg tgg aca gat 1446
 Lys Trp Met Tyr Val Gly Gln Gln His Ala Gly Gln Thr Trp Thr Asp
 435 440 445
 tta act ggc aat cac gcg gcg gtt acg att aat ggt gat ggc tgg 1494
 Leu Thr Gly Asn His Ala Ala Ser Val Thr Ile Asn Gly Asp Gly Trp
 450 455 460
 ggc gaa ttc ttt aca aat gga gga tct gta tcc gtg tat gtg aac caa 1542
 Gly Glu Phe Phe Thr Asn Gly Gly Ser Val Ser Val Tyr Val Asn Gln
 465 470 475 480
 taataaaaag ccttgagaag ggattcctcc ctaactcaag gctttcttta tgcgttttag 1602
 ctcaacgctt ctacgaagct tta 1625

<210>6
 <211>30
 <212>DNA
 <213>Artificial Sequence

<400>6
 atgatgcagt attttgagtg gcatttggaa 30

<210>7
 <211>33
 <212>DNA
 <213>Artificial Sequence

<400>7
 tatgagtggc atttgccaaa cgacgggcag cat 33

<210>8
 <211>33
 <212>DNA
 <213>Artificial Sequence

<400>8
 ccagcctaca aaggtagtag tcaggcggat gtt 33

<210>9
 <211>21
 <212>DNA
 <213>Artificial Sequence

<400>9
 gcacagcttc aacgagctat t 21

5
<210>10
<211>21
<212>DNA
<213>Artificial Sequence

<400>10
tttcgacttt ccagggcgta a 21

10
<210>11
<211>33
<212>DNA
<213>Artificial Sequence

<400>11
catattttcc gctttcaaaa tacgaactgg aac 33

15
<210>12
<211>33
<212>DNA
<213>Artificial Sequence

20
<400>12
aactggcgag tggatgatga gaacggtaat tat 33

<210>13
<211>25
<212>DNA
<213>Artificial Sequence

25
<400>13
tggatgaaga gttcggtaat tatga 25

30
<210>14
<211>33
<212>DNA
<213>Artificial Sequence

<400>14
aatatcgact ttagtcgtcc agaagtacaa gat 33

35
<210>15
<211>33
<212>DNA
<213>Artificial Sequence

40
<400>15
agtcattccag aggtcgtaga tgagttgaag gat 33

<210>16
<211>33
<212>DNA
<213>Artificial Sequence

45
<400>16
gttgactggg atgagcgcaa acaagaaat cat

50
<210>17
<211>34
<212>DNA
<213>Artificial Sequence

55

5
<400>17
atttgccaaa tgacgggcag cattggaatc gggtt 34

<210>18
<211>34
<212>DNA
<213>Artificial Sequence

10
<400>18
aaccgattcc aatgctgccc gtcatttggc aaat 34

<210>19
<211>40
<212>DNA
<213>Artificial Sequence

15
<400>19
gggtcgacca gcacaagccg atggattgaa cggtagcatg 40

<210>20
<211>29
<212>DNA
<213>Artificial Sequence

20
<400>20
taaagctttt gttattgggtt cacgtacac 29

<210>21
<211>30
<212>DNA
<213>Artificial Sequence

25
<400>21
gagtcgacca gcacaagccc atcataatgg 30

<210>22
<211>21
<212>DNA
<213>Artificial Sequence

30
<400>22
taaagcttca atttatattg g 21

<210>23
<211>27
<212>DNA
<213>Artificial Sequence

35
<400>23
ccagatctac ttaccatttt agagtca 27

40

45

50

55

Claims

1. A gene encoding a mutant α -amylase having an amino acid sequence selected from the group consisting of:

(a) amino acid sequences obtained by making replacement or deletion of at least one residue of amino acid residues respectively corresponding to the 11th Tyr, 16th Glu, 49th Asn, 84th Glu, 144th Ser, 167th Gln, 169th Tyr, 178th Ala, 188th Glu, 190th Asn, 205th His and 209th Gln in the amino acid sequence set forth in SEQ ID NO:1 in an α -amylase having said amino acid sequence, or

(b) amino acid sequences obtained by making replacement of a sequence corresponding to 11 to 100 amino acid residues from the amino terminal in the amino acid sequence set forth in SEQ ID NO:1 in an α -amylase having said amino acid sequence by an amino acid sequence of another liquefying α -amylase corresponding to said sequence of the amino acid residues or

(c) amino acid sequences obtained by making replacement or deletion of the above (a) or (b) in the amino acid sequence having a homology of at least 70% to the amino acid sequence set forth in SEQ ID NO:1.

2. The gene according to Claim 1, wherein in the mutant α -amylase a sequence corresponding to amino acid residues from the 1st Asp to the 19th Gly in the amino acid sequence of SEQ ID NO:1 is replaced by an amino acid sequence of another liquefying α -amylase corresponding to said amino acid sequence.

3. The gene according to Claim 1 or 2, wherein said another liquefying α -amylase has the amino acid sequence set forth in SEQ ID NO:2.

4. The gene according to any one of Claims 1 to 3 wherein at least two kinds of replacement or deletion as described above are combined with each other.

5. The gene according to Claim 4, wherein the replacement of the amino acid residue comprises replacing an amino acid residue corresponding to the 11th Tyr in the amino acid sequence of SEQ ID NO:1 by Phe, an amino acid residue corresponding to the 16th Glu by Pro, an amino acid residue corresponding to the 49th Asn by Ser, an amino acid residue corresponding to the 167th Gln by Glu, an amino acid residue corresponding to the 169th Tyr by Lys, an amino acid residue corresponding to the 190th Asn by Phe, an amino acid residue corresponding to the 205th His by Arg, or an amino acid residue corresponding to the 209th Gln by Val, and the replacement of the amino acid sequence comprises replacing an amino acid sequence from the 1st Asp to the 19th Gly in the amino acid sequence of SEQ ID NO:1 by an amino acid sequence from the 1st His to the 21st Gly in the amino acid sequence set forth in SEQ ID NO:2.

6. A vector comprising the gene according to any one of Claims 1 to 5.

7. Cells transformed by the vector according to Claim 6.

8. A process for producing a mutant α -amylase, comprising culturing the transformed cells according to Claim 7.

9. A mutant α -amylase encoded by the gene of any one of Claims 1 to 5 or obtainable by the method of claim 8.

10. A detergent composition comprising the mutant α -amylase according to Claim 9.

Fig. 1




Fig. 2





Fig. 3





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 11 1911

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 99 23211 A (NOVONORDISK AS) 14 May 1999 (1999-05-14) * abstract * * page 45 * * page 51 * * page 52 * * page 54 *	1,4-10	C12N15/56 C12N9/28 C12N1/21 C11D3/386 //(C12N1/21, C12R1:125)
A	CONRAD BIRGIT ET AL: "Hybrid Bacillus amyloliquefaciens X Bacillus licheniformis alpha-amylases: Construction, properties and sequence determinants." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 230, no. 2, 1995, pages 481-490, XP000960424 ISSN: 0014-2956 * abstract; figure 2 *	1-5	
A	IGARASHI KAZUAKI ET AL: "Improved thermostability of a Bacillus alpha-amylase by deletion of an arginine-glycine residue is caused by enhanced calcium binding." BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 248, no. 2, 20 July 1998 (1998-07-20), pages 372-377, XP002901159 ISSN: 0006-291X * the whole document *		TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N C11D
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 8 November 2000	Examiner Lejeune, R
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1603 03.02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 11 1911

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-11-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9923211 A	14-05-1999	AU 9737398 A	24-05-1999
		BR 9813328 A	22-08-2000
		EP 1027428 A	16-08-2000
<hr/>			

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82